



Deborah A. Faryniarz, M.D., Inc.

*Sports Medicine and Shoulder Fellowship,
The Hospital for Special Surgery,
FORUM, Society of Fellowship Trained
Women in Orthopaedic Surgery*

Grady L. Jeter, M.D., Inc.

*Member Arthroscopy Association of North America,
International Society of Arthroscopy,
Knee Surgery and Orthopaedic Sports Medicine,
Founding Member Ikeuchi International Society for Arthroscopy and
Musculoskeletal Endoscopy*

Martin Trieb, M.D., Inc.

Member Arthroscopy Association of North America

*Diplomates American Board of Orthopaedic Surgery • Members American Orthopaedic Society for Sports Medicine
Fellows American Academy of Orthopaedic Surgeons*

Postoperative Instructions Rotator Cuff Repair

Immobilizer: Should be worn at all times (including sleeping) except for hygiene and therapeutic exercises until approved by Dr. Faryniarz (usually 4-6 weeks).

Ice

- Apply an ice pack or Game Ready icing machine to the shoulder as often as possible for the first 3-7 days to reduce pain and swelling. You may ice as often as 30 minutes every hour but not less than 3x/day. If using Game Ready, use the no pressure or low pressure setting.

Medications

- Pain medication is injected into the joint during surgery and will wear off within 8-12 hrs.
- Most patients require a narcotic (i.e. Percocet or Vicodin) for a short period of time. Please follow the prescription directions.
- Common side effects of pain medications include drowsiness, nausea, and constipation. To minimize side effects, take medications with food. You may take an over-the-counter laxative (i.e. Colace) if needed. Do not drive or operate any machinery while on narcotics.
- You may take anti-inflammatory medication (400 mg Motrin/Advil) between doses of narcotics to help control pain. Take with food to avoid stomach upset. Do not take if you have a history of ulcers.
- If you experiencing any problems or uncomfortable side effects, please contact our office.

Wound Care

- Remove the dressing, but not the Steri-strips, two days after surgery. You may cover the wound with Band-aids if any drainage is present. It is normal to have some drainage and swelling after surgery. If the dressing becomes soaked, you may reinforce with additional dressing.
- Avoid any ointments around the incisions.
- Keep wounds clean and dry for the first 5 days. If you wish to shower during this time, cover your shoulder/sling with a large plastic bag or take a sponge bath. After 5 days, you may shower, but do not soak in a bathtub or hot tub until after the first post-op visit.
- Your post-op appointment is on your surgery instruction sheet (usually 7-14 days after your surgery). If you need to reschedule, please call my assistant at (408) 559-3888.
- **If you develop a fever (temperature > 101.5), excessive drainage, redness around the incision, increased pain or numbness in your arm, or difficulty breathing notify Dr. Faryniarz at (408) 559-3888.**

Activity

- You should start PT a few days after your surgery. Please call our office if have not received an appointment or PT script. No active elbow bending if biceps tenodesis done.
- You may be more comfortable sleeping in a recliner with a pillow behind your arm and under your forearm for support.
- You may take your arm out of the sling several times a day to passively range your elbow. If you develop pain in your elbow or numbness in your ring or pinky finger, take your arm out of the sling and straighten your elbow for several hours. If persists, call Dr. Faryniarz.
- **No driving is allowed until instructed by Dr. Faryniarz (usually 4-6 weeks).**
- If pain permits, you may return to (sedentary) work or school 3-4 days after surgery.
- Return to work with lifting or overhead reaching will be determined by Dr. Faryniarz.